

**A novel and highly specific phage endolysin  
cell wall binding domain for detection of  
*Bacillus cereus***

*Minsuk Kong,<sup>†,&</sup> Jieun Sim,<sup>‡,||,#,&</sup> Taejoon Kang,<sup>‡,||</sup> Hoang Hiep Nguyen,<sup>‡,#</sup> Hyun Kyu Park,<sup>⊥</sup>  
Bong Hyun Chung,<sup>‡,||,#,\*</sup> and Sangryeol Ryu<sup>‡,\*</sup>*

<sup>†</sup>Department of Food and Animal Biotechnology, Research Institute of Agriculture and Life Sciences, and Center for Food and Bioconvergence, Seoul National University, Seoul 151-742, Korea


<sup>‡</sup>BioNanotechnology Research Center and <sup>||</sup>BioNano Health Guard Research Center, KRIBB, Daejeon 305-806, Korea

<sup>#</sup>Major of Nanobiotechnology, UST, Daejeon 305-806, Korea

<sup>⊥</sup>MicoBioMed Ltd, Daejeon 305-806, Korea

**Corresponding Author**

\*E-mail: sangryu@snu.ac.kr (S.R.); chungbh@kribb.re.kr (B.H.C.)



**Fig. S1.** Optical and fluorescent images of various bacterial species after the addition of GFP-LysPBC1\_CBD.