

Supporting Information

Diagnosis of Tamiflu-Resistant Influenza Virus in Human Nasal Fluid and Saliva using Surface-enhanced Raman Scattering

Gayoung Eom^{a,b}, Ahreum Hwang^{a,b}, Hongki Kim^a, Siyeong Yang^b, Do Kyung Lee^c,
Sinae Song^a, Kab Ha^a, Jinyoung Jeong^{d,e}, Juyeon Jung^{a,d}, Eun-Kyung Lim^{a,d,*},
and Taejoon Kang^{a,d,*}

^aBionanotechnology Research Center, KRIBB, Daejeon 34141, Republic of Korea

^bDepartment of Chemistry, KAIST, Daejeon 34141, Republic of Korea

^cBioNano Health Guard Research Center, KRIBB, Daejeon 34141, Republic of Korea

^dDepartment of Nanobiotechnology, KRIBB School of Biotechnology, UST, Daejeon 34113,
Republic of Korea

^eEnvironmental Disease Research Center, KRIBB, Daejeon 34141, Republic of Korea

*E-mail: kangtaejoon@kribb.re.kr (T. Kang); eklim1112@kribb.re.kr (E.-K. Lim)




Figure S1. Plot of absorbance values depending on the type of influenza virus (WT virus, mutant virus, and control).




Figure S2. Extinction spectrum of urchin Au NPs.




Figure S3. Full SERS spectra corresponding to Figure 3b.




Figure S4. SEM images of OHT-functionalized glass slides after treatment of nasal fluid and saliva samples with WT virus or no virus.